



Clement, M., Pearson, J., Gras, S., van den Berg, H. A., Lissina, A., Llewellyn-Lacey, S., Willis, M., Dockree, T., McLaren, J. E., Ekeruche-Makinde, J., Gostick, E., Robertson, N. P., Rossjohn, J., Burrows, S. R., Price, D. A., Wong, S., Peakman, M., Skowera, A., & Wooldridge, L. (2016). Targeted suppression of autoreactive CD8⁺ T-cell activation using blocking anti-CD8 antibodies. *Scientific Reports*, 6, [35332]. <https://doi.org/10.1038/srep35332>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1038/srep35332](https://doi.org/10.1038/srep35332)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Nature at <http://www.nature.com/articles/srep35332>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Targeted suppression of autoreactive CD8⁺ T-cell activation using blocking anti-CD8 antibodies

Mathew Clement, James A. Pearson, Stephanie Gras, Hugo A. van den Berg, Anya Lissina, Sian Llewellyn-Lacey, Mark D. Willis, Tamsin Dockree, James E. McLaren, Julia Ekeruche-Makinde, Emma Gostick, Neil P. Robertson, Jamie Rossjohn, Scott R. Burrows, David A. Price, F. Susan Wong, Mark Peakman, Ania Skowera, Linda Wooldridge

Figure S1




Figure S1: Affinities and kinetics of GILGFVFTL-HLA-A*0201 binding to the ALF3 and ALF8 TCRs. Surface plasmon resonance binding curves of the ALF3 and ALF8 TCRs in solid phase tested against the GILGFVFTL-HLA-A*0201-M1 complex in fluid phase across a range of concentrations (0.78–200 µM). The mean \pm SEM of two replicate assays is shown.

Figure S2




Figure S2: Combinatorial peptide library screen of the 1E6 CD8⁺ T-cell clone in the absence of an intact pMHC/CD8 interaction. 6×10^4 C1R-A*0201 D227K/T228A target cells were pulsed in duplicate with each mixture from a 10mer combinatorial peptide library (100 μ M) for 2 hours at 37°C. 3×10^4 clonal 1E6 CD8⁺ T-cells were then added and the plates were incubated overnight at 37°C. Supernatants were assayed for MIP-1 β by ELISA. The index peptide sequence is shown as red bars in each position.

CD8 ⁺ T-cell clone ID	No treatment (pEC ₅₀ MIP-1 β)	0.25 μ g/ml DK25 (pEC ₅₀ MIP-1 β)	0.5 μ g/ml DK25 (pEC ₅₀ MIP-1 β)	1 μ g/ml DK25 (pEC ₅₀ MIP-1 β)
1E6	6.74921	-	-	-
3F2	6.02653	-	-	-
4C6	6.34537	-	-	-
ALF3	8.27954	7.9456	7.64974	7.76992
SB27	8.82749	8.81323	8.55271	8.72455
MCNLV	9.40681	8.82467	8.7935	8.60756

Table S1A: Functional sensitivity, measured by MIP-1 β production, expressed as pEC₅₀ (Figure 5)¹.

CD8 ⁺ T-cell clone ID	No treatment (pEC ₅₀ lysis)	0.25 μ g/ml DK25 (pEC ₅₀ lysis)	0.5 μ g/ml DK25 (pEC ₅₀ lysis)	1 μ g/ml DK25 (pEC ₅₀ lysis)
1E6	7.34425	-	-	-
3F2	7.06595	-	-	-
4C6	7.58755	-	-	-
ALF3	9.41053	8.69299	8.55927	8.62777
SB27	9.22728	9.06544	8.90326	8.44131
MCNLV	9.29825	9.11505	8.51186	8.24729

Table S1B: Functional sensitivity, measured by killing activity, expressed as pEC₅₀ (Figure 6)¹.

CD8 ⁺ T-cell clone ID	0.25 μ g/ml DK25 (delta pEC ₅₀ MIP-1 β)	0.5 μ g/ml DK25 (delta pEC ₅₀ MIP-1 β)	1 μ g/ml DK25 (delta pEC ₅₀ MIP-1 β)
1E6	>1.75	>1.75	>1.75
3F2	>1.03	>1.03	>1.03
4C6	>1.35	>1.35	>1.35
ALF3	0.33394	0.6298	0.50962
SB27	0.01426	0.27478	0.10299
MCNLV	0.58214	0.61331	0.79925

Table S1C: DK25-induced shifts in functional sensitivity, measured by MIP-1 β production, expressed as delta pEC₅₀ (Figure 5).

CD8 ⁺ T-cell clone ID	0.25 μ g/ml DK25 (delta pEC ₅₀ lysis)	0.5 μ g/ml DK25 (delta pEC ₅₀ lysis)	1 μ g/ml DK25 (delta pEC ₅₀ lysis)
1E6	>2.34	>2.34	>2.34
3F2	>2.07	>2.07	>2.07
4C6	>2.59	>2.59	>2.59
ALF3	0.71754	0.85126	0.78276
SB27	0.16184	0.32402	0.78597
MCNLV	0.78639	0.78639	1.05096

Table S1D: DK25-induced shifts in functional sensitivity, measured by killing activity, expressed as delta pEC₅₀ (Figure 6).

1. Sprent P, Smeeton NC. Applied nonparametric statistical methods. *Chapman & Hall/CRC, London* (2007).